

LIPSCHITZ CONDITIONS AND THE DISTANCE RATIO METRIC

SLAVKO SIMIĆ AND MATTI VUORINEN

ABSTRACT. We give study the Lipschitz continuity of Möbius transformations of a punctured disk onto another punctured disk with respect to the distance ratio metric.

1. INTRODUCTION

During the past thirty years the theory of quasiconformal maps has been studied in various contexts such as in Euclidean, Banach, or even metric spaces. It has turned out that while some classical tools based on conformal invariants, real analysis and measure theory are no longer useful beyond the Euclidean context, the notion of a metric space and related notions still provide a useful conceptual framework. This has led to the study of the geometry defined by various metrics and to the key role of metrics in recent theory of quasiconformality. See e.g. [CCQ, HIMPS, HPWW, K, RT1, RT2, V].

Distance ratio metric. One of these metrics is the distance ratio metric. For a subdomain $G \subset \mathbb{R}^n$ and for $x, y \in G$ the *distance ratio metric* j_G is defined by

$$(1.1) \quad j_G(x, y) = \log \left(1 + \frac{|x - y|}{\min\{d_G(x), d_G(y)\}} \right),$$

where $d_G(x)$ denotes the Euclidean distance from x to ∂G . If $G_1 \subset G$ is a proper subdomain then for $x, y \in G_1$ clearly

$$(1.2) \quad j_G(x, y) \leq j_{G_1}(x, y).$$

Moreover, the numerical value of the metric is highly sensitive to boundary variation, the left and right sides of (1.2) are not comparable even if $G_1 = G \setminus \{p\}, p \in G$.

The distance ratio metric was introduced by F.W. Gehring and B.P. Palka [GP] and in the above, simplified, form by M. Vuorinen [Vu1] and it is frequently used in the study of hyperbolic type metrics [HIMPS] and geometric theory of functions. It is a basic fact that the above j -metric is closely related to the hyperbolic metric both for the unit ball $\mathbb{B}^n \subset \mathbb{R}^n$ and for the Poincaré half-space \mathbb{H}^n , [Vu2].

Quasi-invariance of j_G .

Given domains $G, G' \subset \mathbb{R}^n$ and an open continuous mapping $f : G \rightarrow G'$ with $fG \subset G'$ we consider the following condition: there exists a constant $C \geq 1$ such that for all $x, y \in G$ we have

$$(1.3) \quad j_{G'}(f(x), f(y)) \leq C j_G(x, y),$$

or, equivalently, that the mapping

$$f : (G, j_G) \rightarrow (G', j_{G'})$$

between metric spaces is Lipschitz continuous with the Lipschitz constant C .

The hyperbolic metric in the unit ball or half space are Möbius invariant. However, the distance ratio metric j_G is not invariant under Möbius transformations. Therefore, it is natural to ask what the Lipschitz constants are for these metrics under conformal mappings or Möbius transformations in higher dimension. F.W. Gehring and B.G. Osgood proved that these metrics are not changed by more than a factor 2 under Möbius transformations, see [GO, proof of Theorem 4]:

Theorem 1.4. *If D and D' are proper subdomains of \mathbb{R}^n and if f is a Möbius transformation of D onto D' , then for all $x, y \in D$*

$$\frac{1}{2}j_D(x, y) \leq j_{D'}(f(x), f(y)) \leq 2j_D(x, y).$$

It is easy to see that for a Möbius transformation $f : \mathbb{B}^n \rightarrow \mathbb{B}^n$ with $f(0) \neq 0$, and $x, y \in \mathbb{B}^n, x \neq y$, the $j_{\mathbb{B}^n}$ distances need not be the same. On the other hand, the next theorem from [SVW], conjectured in [KVZ], yields a sharp form of Theorem 1.4 for Möbius automorphisms of the unit ball.

Theorem 1.5. *A Möbius transformation $f : \mathbb{B}^n \rightarrow \mathbb{B}^n = f(\mathbb{B}^n)$ satisfies*

$$j_{\mathbb{B}^n}(f(x), f(y)) \leq (1 + |f(0)|)j_{\mathbb{B}^n}(x, y)$$

for all $x, y \in \mathbb{B}^n$. The constant is best possible.

A similar result for a punctured disk was conjectured in [SVW]. The next theorem, our main result, settles this conjecture from [SVW] in the affirmative.

Theorem 1.6. *Let $a \in \mathbb{B}^2$ and $h : \mathbb{B}^2 \setminus \{0\} \rightarrow \mathbb{B}^2 \setminus \{a\}$ be a Möbius transformation with $h(0) = a$. Then for $x, y \in \mathbb{B}^2 \setminus \{0\}$*

$$j_{\mathbb{B}^2 \setminus \{a\}}(h(x), h(y)) \leq C(a)j_{\mathbb{B}^2 \setminus \{0\}}(x, y),$$

where the constant $C(a) = 1 + (\log \frac{2+|a|}{2-|a|})/\log 3$ is best possible.

Clearly the constant $C(a) < 1 + |a| < 2$ for all $a \in \mathbb{B}^2$ and hence the constant in Theorem 1.6 is smaller than the constant $1 + |f(0)|$ in Theorem 1.5 and far smaller than the constant 2 in Theorem 1.4.

If $a = 0$ in Theorem 1.6, then h is a rotation of the unit disk and hence a Euclidean isometry. Note that $C(0) = 1$, i.e. the result is sharp in this case.

The proof is based on Theorem 2.1 below and on Lemma 2.4, a monotone form of l'Hôpital's rule from [AVV, Theorem 1.25].

2. PRELIMINARY RESULTS

In view of the definition of the distance ratio metric it is natural to expect that some properties of the logarithm will be needed. In the earlier paper [SVW], the classical Bernoulli inequality [Vu2, (3.6)] was applied for this purpose. Apparently now some other inequalities are needed and we use the following result, which is precise and allows us to get rid of logarithms in further calculations.

Theorem 2.1. Let D and D' be proper subdomains of \mathbb{R}^n . For an open continuous mapping $f : D \rightarrow D'$ denote

$$X = X(z, w) := \frac{|z - w|}{\min\{d_D(z), d_D(w)\}}; \quad Y = Y(z, w) := \frac{|z - w|}{|f(z) - f(w)|} \frac{\min\{d_{D'}(f(z)), d_{D'}(f(w))\}}{\min\{d_D(z), d_D(w)\}}.$$

If there exists q , $0 \leq q \leq 1$ such that

$$(2.2) \quad q \leq Y + \frac{Y - 1}{X + 1},$$

then the inequality

$$j_{D'}(f(z), f(w)) \leq \frac{2}{1 + q} j_D(z, w),$$

holds for all $z, w \in D$.

Proof. The proof is based on the following assertion.

Lemma 2.3. For $a \geq 0, q \in [0, 1]$, we have

$$\log\left(\frac{q + e^a}{1 + qe^a}\right) \leq \frac{1 - q}{1 + q}a.$$

Proof. Denote

$$f(a, q) := \log\left(\frac{q + e^a}{1 + qe^a}\right) - \frac{1 - q}{1 + q}a.$$

By differentiation, we have

$$f'_a(a, q) = -\frac{q(1 - q)}{1 + q} \frac{(e^a - 1)^2}{(1 + qe^a)(q + e^a)},$$

we conclude that

$$f(a, q) \leq f(0, q) = 0.$$

□

Now, since

$$X = \frac{|z - w|}{\min\{d_D(z), d_D(w)\}} = \exp(j_D(z, w)) - 1,$$

and

$$Y = \frac{|z - w|}{|f(z) - f(w)|} \frac{\min\{d_{D'}(f(z)), d_{D'}(f(w))\}}{\min\{d_D(z), d_D(w)\}} = \frac{\exp(j_D(z, w)) - 1}{\exp(j_{D'}(f(z), f(w))) - 1},$$

the condition (2.2) is equivalent to

$$\exp(j_{D'}(f(z), f(w))) \leq \exp(j_D(z, w)) \left(\frac{q + e^{j_D(z, w)}}{1 + qe^{j_D(z, w)}} \right).$$

Therefore, by Lemma 2.3, we get

$$\begin{aligned} j_{D'}(f(z), f(w)) &\leq j_D(z, w) + \log\left(\frac{q + e^{j_D(z, w)}}{1 + qe^{j_D(z, w)}}\right) \\ &\leq j_D(z, w) + \frac{1 - q}{1 + q}j_D(z, w) = \frac{2}{1 + q}j_D(z, w). \end{aligned}$$

□

In the sequel we shall need the so-called *monotone form of l'Hôpital's rule*.

Lemma 2.4. [AVV, Theorem 1.25]. For $-\infty < a < b < \infty$, let $f, g : [a, b] \rightarrow \mathbb{R}$ be continuous on $[a, b]$, and be differentiable on (a, b) , and let $g'(x) \neq 0$ on (a, b) . If $f'(x)/g'(x)$ is increasing(decreasing) on (a, b) , then so are

$$\frac{f(x) - f(a)}{g(x) - g(a)} \quad \text{and} \quad \frac{f(x) - f(b)}{g(x) - g(b)}.$$

If $f'(x)/g'(x)$ is strictly monotone, then the monotonicity in the conclusion is also strict.

Lemma 2.4 has found numerous applications recently. See the bibliography of [AVZ] for a long list of applications to inequalities.

Lemma 2.5. For positive numbers A, B, D and $0 < C < 1, \theta \geq 0$, we have

1. The inequality

$$1 + \frac{B}{D}\theta(1 + \frac{D}{1+A})(1 + \frac{B}{1-C}\theta) \leq (1 + \frac{B}{D}\theta)(1 + \frac{B}{1-C}\theta),$$

holds if and only if $B\theta \leq A + C$;

2. The function

$$\frac{\log(1 + \frac{B}{1-C}\theta)}{\log(1 + \frac{B}{D}\theta)}$$

is monotone increasing (decreasing) in θ if $C + D < 1$ ($C + D > 1$).

Proof. Proof of the first part follows by direct calculation.

For the second part, set

$$f_1(\theta) = \log(1 + \frac{B}{1-C}\theta), f_1(0) = 0; \quad f_2(\theta) = \log(1 + \frac{B}{D}\theta), f_2(0) = 0.$$

Since

$$\frac{f_1'(\theta)}{f_2'(\theta)} = \frac{D + B\theta}{1 - C + B\theta} = 1 + \frac{C + D - 1}{1 - C + B\theta},$$

the proof follows according to Lemma 2.4. □

3. PROOF OF THEOREM 1.6

3.1. Proof of Theorem 1.6.

For the proof, define $h(z) = \frac{z+a}{1+\bar{a}z}$ and suppose in the sequel that $|z| \geq |w|$. Then

$$j_G(z, w) = \log\left(1 + \frac{|z - w|}{\min\{|z|, |w|, 1 - |z|, 1 - |w|\}}\right) = \log\left(1 + \frac{|z - w|}{\min\{|w|, 1 - |z|\}}\right),$$

and

$$j_{G'}(h(z), h(w)) = \log\left(1 + \frac{|h(z) - h(w)|}{T}\right),$$

where

$$T = T(a, z, w) := \min\{|h(z) - a|, |h(w) - a|, 1 - |h(z)|, 1 - |h(w)|\}.$$

In concert with the definition of the number T , the proof is divided into four cases. We shall consider each case separately applying Bernoulli inequality in the first case, its stronger form from Theorem 2.1 in the second one and a direct approach in the last two cases.

$$1. \quad T = |h(z) - a|.$$

Since $|h(z) - a| = \frac{(1-|a|^2)|z|}{|1+\bar{a}z|}$ and $|h(z) - h(w)| = \frac{(1-|a|^2)|z-w|}{|1+\bar{a}z||1+\bar{a}w|}$, we have

$$j_{G'}(h(z), h(w)) = \log\left(1 + \frac{|z-w|}{|z||1+\bar{a}w|}\right).$$

Suppose firstly that $|w| \leq 1 - |z|$. Since also $|w| \leq 1 - |z| \leq 1 - |w|$, we conclude that $0 \leq |w| \leq 1/2$. Hence, by the Bernoulli inequality (see e.g. [Vu2, (3.6)]), we get

$$\begin{aligned} j_{G'}(h(z), h(w)) &\leq \log\left(1 + \frac{|z-w|}{|z|(1-|a||w|)}\right) \leq \log\left(1 + \frac{|z-w|}{|w|(1-\frac{|a|}{2})}\right) \\ &\leq \frac{1}{1-\frac{|a|}{2}} \log\left(1 + \frac{|z-w|}{|w|}\right) = \frac{1}{1-\frac{|a|}{2}} j_G(z, w). \end{aligned}$$

Suppose now $1 - |z| \leq |w| (\leq |z|)$. Then $1/2 \leq |z| < 1$.

Since in this case $(|z| - \frac{1}{2})(2 - |a|(1 + |z|)) \geq 0$, we easily obtain that

$$\frac{1}{|z|(1-|a||z|)} \leq \frac{1}{(1-\frac{|a|}{2})(1-|z|)}.$$

Hence,

$$\begin{aligned} j_{G'}(h(z), h(w)) &\leq \log\left(1 + \frac{|z-w|}{|z|(1-|a||w|)}\right) \leq \log\left(1 + \frac{|z-w|}{|z|(1-|a||z|)}\right) \\ &\leq \log\left(1 + \frac{|z-w|}{(1-\frac{|a|}{2})(1-|z|)}\right) \leq \frac{1}{1-\frac{|a|}{2}} \log\left(1 + \frac{|z-w|}{1-|z|}\right) = \frac{1}{1-\frac{|a|}{2}} j_G(z, w). \end{aligned}$$

$$2. \quad T = |h(w) - a|.$$

This case can be treated by means of Theorem 2.1 with the same resulting constant $C_1(a) = \frac{2}{2-|a|}$.

Indeed, in terms of Theorem 2.1, we consider firstly the case $|w| \leq 1 - |z|$.

We get

$$X = \frac{|z-w|}{|w|} \geq \frac{|z|-|w|}{|w|} = \frac{|z|}{|w|} - 1 = X^*,$$

and

$$Y = |1 + \bar{a}z| \geq 1 - |a||z| = Y^*.$$

Therefore,

$$\begin{aligned} Y + \frac{Y-1}{X+1} &\geq Y^* - \frac{1-Y^*}{1+X^*} = 1 - |a||z| - |a||z| \frac{|w|}{|z|} \\ &= 1 - |a|(|w| + |z|) \geq 1 - |a| = q. \end{aligned}$$

In the second case, i.e. when $1 - |z| \leq |w|$, we want to show that

$$Y + \frac{Y-1}{X+1} \geq 1 - |a|, \text{ with } X = \frac{|z-w|}{1-|z|}, Y = |1 + \bar{a}z| \frac{|w|}{1-|z|}.$$

This is equivalent to

$$(Y - (1 - |a|))(1 + X) + Y \geq 1.$$

Since in this case

$$X = \frac{|z-w|}{1-|z|} \geq \frac{|z|-|w|}{1-|z|} := X^*$$

and

$$Y = |1 + \bar{a}z| \frac{|w|}{1-|z|} \geq (1 - |a||z|) \frac{|w|}{1-|z|} = 1 - |a||z| + (|w| + |z| - 1) \frac{1 - |a||z|}{1-|z|} := Y^*,$$

we get

$$\begin{aligned} & (Y - (1 - |a|))(1 + X) + Y - 1 \geq (Y^* - (1 - |a|))(1 + X^*) + Y^* - 1 \\ &= \left[|a|(1 - |z|) + (|w| + |z| - 1) \frac{1 - |a||z|}{1-|z|} \right] \frac{1 - |w|}{1-|z|} - |a||z| + (|w| + |z| - 1) \frac{1 - |a||z|}{1-|z|} \\ &\geq |a|(1 - |w| - |z|) + (|w| + |z| - 1) \frac{1 - |a||z|}{1-|z|} = (|w| + |z| - 1) \frac{1 - |a|}{1-|z|} \geq 0. \end{aligned}$$

Therefore by Theorem 2.1, in both cases we get

$$j_{G'}(h(z), h(w)) \leq \frac{2}{1+q} j_G(z, w) = \frac{2}{2-|a|} j_G(z, w) = C_1(a) j_G(z, w).$$

3. $T = 1 - |h(z)|$.

In this case, applying well-known assertions

$$|1 + \bar{a}z|^2 - |a + z|^2 = (1 - |a|^2)(1 - |z|^2); \quad |h(z)| \leq \frac{|a| + |z|}{1 + |a||z|},$$

and

$$|1 + \bar{a}w| \geq 1 - |a||w| (\geq 1 - |a||z|),$$

we get

$$\begin{aligned} j_{G'}(h(z), h(w)) &= \log \left(1 + \frac{|z-w|(1-|a|^2)}{|1 + \bar{a}w|(|1 + \bar{a}z| - |z+a|)} \right) = \log \left(1 + \frac{|z-w|(|1 + \bar{a}z| + |z+a|)}{|1 + \bar{a}w|(1-|z|^2)} \right) \\ &= \log \left(1 + \frac{|z-w|}{1-|z|^2} \left| 1 + \frac{\bar{a}(z-w)}{1 + \bar{a}w} \right| \left(1 + \frac{|z+a|}{|1 + \bar{a}z|} \right) \right) \leq \log \left(1 + \frac{|z-w|}{1-|z|} \left(1 + \frac{|a||z-w|}{1-|a||w|} \right) \left(1 + \frac{|a|(1-|z|)}{1+|a||z|} \right) \right). \end{aligned}$$

Applying here Lemma 2.5, part 1., with

$$A = |a||z|, \quad B = |a|, \quad C = |a||w|, \quad D = |a|(1-|z|), \quad \theta = |z-w|,$$

we obtain

$$(3.2) \quad j_{G'}(h(z), h(w)) \leq \log \left[\left(1 + \frac{|z-w|}{1-|z|} \right) \left(1 + \frac{|a||z-w|}{1-|a||w|} \right) \right].$$

Suppose that $1 - |z| \leq |w| (\leq |z|)$. By Lemma 2.5, part 2., with

$$B = |a|, \quad C = |a||z|, \quad D = |a|(1 - |z|), \quad \theta = |z - w|,$$

we get

$$\begin{aligned} J(z, w; a) &:= \frac{j_{G'}(h(z), h(w))}{j_G(z, w)} \leq 1 + \frac{\log(1 + \frac{|a||z-w|}{1-|a||z|})}{\log(1 + \frac{|z-w|}{1-|z|})} \\ &\leq 1 + \frac{\log(1 + \frac{2|a||z|}{1-|a||z|})}{\log(1 + \frac{2|z|}{1-|z|})}, \end{aligned}$$

because in this case we have $C + D = |a| < 1$ and $|z - w| \leq 2|z|$.

Since the last function is monotone decreasing in $|z|$ and $|z| \geq 1/2$, we obtain

$$J(z, w; a) \leq 1 + \frac{\log(\frac{1+\frac{1}{2}|a|}{1-\frac{1}{2}|a|})}{\log(\frac{1+\frac{1}{2}}{1-\frac{1}{2}})} = 1 + (\log \frac{2+|a|}{2-|a|}) / \log 3 := C_2(a).$$

Let now $|w| \leq 1 - |z| (\leq 1 - |w|)$. The estimation (3.2) and Lemma 2.5, part 2., with

$$B = |a|, \quad C = D = |a||w|, \quad \theta = |z - w|,$$

yield

$$\begin{aligned} J(z, w; a) &\leq \frac{\log\left[\left(1 + \frac{|z-w|}{1-|z|}\right)\left(1 + \frac{|a||z-w|}{1-|a||w|}\right)\right]}{\log\left(1 + \frac{|z-w|}{|w|}\right)} \leq \frac{\log\left[\left(1 + \frac{|z-w|}{|w|}\right)\left(1 + \frac{|a||z-w|}{1-|a||w|}\right)\right]}{\log\left(1 + \frac{|z-w|}{|w|}\right)} \\ &= 1 + \frac{\log\left(1 + \frac{|a||z-w|}{1-|a||w|}\right)}{\log\left(1 + \frac{|z-w|}{|w|}\right)} \leq 1 + \frac{\log\left(1 + \frac{|a|}{1-|a||w|}\right)}{\log\left(1 + \frac{1}{|w|}\right)}, \end{aligned}$$

since $C + D = 2|a||w| \leq |a| < 1$ and $0 \leq |z - w| \leq |z| + |w| \leq 1$.

Denote the last function as $g(|w|)$ and let $|w| = r$, $0 < r \leq 1/2$. Since

$$g'(r) = \frac{|a|^2}{(1-r|a|)(1+(1-r)|a|)\log(1+1/r)} + \frac{\log\left(1 + \frac{|a|}{1-|a|r}\right)}{r(1+r)\log^2(1+1/r)} > 0,$$

we finally obtain

$$J(z, w; a) \leq 1 + \frac{\log\left(1 + \frac{|a|}{1-|a|/2}\right)}{\log(1+2)} = C_2(a).$$

4. $T = 1 - |h(w)|$.

This case can be considered analogously with the previous one.

$$j_{G'}(h(z), h(w)) = \log\left(1 + \frac{|z-w|(1-|a|^2)}{|1+\bar{a}z|(|1+\bar{a}w| - |w+a|)}\right) = \log\left(1 + \frac{|z-w|(|1+\bar{a}w| + |w+a|)}{|1+\bar{a}z|(1-|w|^2)}\right)$$

$$= \log\left(1 + \frac{|z-w|}{1-|w|^2} \left|1 + \frac{\bar{a}(\textcolor{red}{w-z})}{1+\bar{a}z}\right| \left(1 + \frac{|w+a|}{|1+\bar{a}w|}\right)\right) \leq \log\left(1 + \frac{|z-w|}{1-|w|} \left(1 + \frac{|a||z-w|}{1-|a||z|}\right) \left(1 + \frac{|a|(1-|w|)}{1+|a||w|}\right)\right)$$

Applying Lemma 2.5, part 1., with

$$A = |a||w|, \quad B = |a|, \quad C = |a||z|, \quad D = |a|(1-|w|), \quad \theta = |z-w|,$$

we obtain

$$(3.3) \quad j_{G'}(h(z), h(w)) \leq \log\left[\left(1 + \frac{|z-w|}{1-|w|}\right) \left(1 + \frac{|a||z-w|}{1-|a||z|}\right)\right].$$

Suppose that $1-|z| \leq |w| (\leq |z|)$. We get

$$\begin{aligned} J(z, w; a) &:= \frac{j_{G'}(h(z), h(w))}{j_G(z, w)} \leq \frac{\log\left[\left(1 + \frac{|z-w|}{1-|z|}\right) \left(1 + \frac{|a||z-w|}{1-|a||z|}\right)\right]}{\log\left(1 + \frac{|z-w|}{1-|z|}\right)} \\ &= 1 + \frac{\log\left(1 + \frac{|a||z-w|}{1-|a||z|}\right)}{\log\left(1 + \frac{|z-w|}{1-|z|}\right)} \end{aligned}$$

and this inequality is already considered above.

In the case $|w| \leq 1-|z| \leq 1-|w|$, we have

$$\begin{aligned} J(z, w; a) &\leq \frac{\log\left[\left(1 + \frac{|z-w|}{1-|w|}\right) \left(1 + \frac{|a||z-w|}{1-|a||z|}\right)\right]}{\log\left(1 + \frac{|z-w|}{|w|}\right)} \leq \frac{\log\left[\left(1 + \frac{|z-w|}{|w|}\right) \left(1 + \frac{|a||z-w|}{1-|a|(1-|w|)}\right)\right]}{\log\left(1 + \frac{|z-w|}{|w|}\right)} \\ &= 1 + \frac{\log\left(1 + \frac{|a||z-w|}{1-|a|(1-|w|)}\right)}{\log\left(1 + \frac{|z-w|}{|w|}\right)} \leq 1 + \frac{\log\left(1 + \frac{|a|}{1-|a|(1-|w|)}\right)}{\log\left(1 + \frac{1}{|w|}\right)}, \end{aligned}$$

where the last inequality follows from Lemma 2.5, part 2., with

$$B = |a|, \quad C = |a|(1-|w|), \quad D = |a||w|, \quad \theta = |z-w|,$$

since $C + D = |a| < 1$ and $|z-w| \leq |z| + |w| \leq 1$.

Denote now $|w| = r$ and let $k(r) = k_1(r)/k_2(r)$ with

$$k_1(r) = \log\left(1 + \frac{|a|}{1-|a|(1-r)}\right); \quad k_2(r) = \log\left(1 + \frac{1}{r}\right).$$

We shall show now that the function $k(r)$ is monotone increasing on the positive part of real axis.

Indeed, since $k_1(\infty) = k_2(\infty) = 0$ and

$$\begin{aligned} k'_1(r)/k'_2(r) &= \frac{|a|^2 r(1+r)}{(1+|a|r)(1-|a|+|a|r)} = \frac{|a|(1+r)}{1+|a|r} \frac{|a|r}{1-|a|+|a|r} \\ &= \left(1 - \frac{1-|a|}{1+|a|r}\right) \left(1 - \frac{1-|a|}{1-|a|+|a|r}\right), \end{aligned}$$

with both functions in parenthesis evidently increasing on \mathbb{R}^+ , the conclusion follows from Lemma 2.4.

Since in this case $0 < r \leq 1/2$, we also obtain that

$$J(z, w; a) \leq 1 + \frac{\log\left(1 + \frac{|a|}{1-|a|/2}\right)}{\log(1+2)} = C_2(a).$$

The constant $C_2(a)$ is sharp since $J(\frac{a}{2|a|}, \frac{-a}{2|a|}; a) = C_2(a)$.

Because $C_2(a) = 1 + (\log \frac{2+a}{2-a}) / \log 3 > \frac{1}{1-\frac{|a|}{2}} = C_1(a)$, we conclude that the best possible upper bound C is $C = C_2(a)$ \square .

Finally, in order to widen the topic started with Conjecture 1, we consider the following:

Let $h : \mathbb{D} \rightarrow \mathbb{D}$ be Möbius map with $h(0) = a$. A challenging problem is to determine best possible j -Lip constants $C(m, a)$ such that

$$j(h^m(z), h^m(w)) \leq C(m, a)j(z, w),$$

for all $z, w \in \mathbb{D}$ and $m \in \mathbb{N}$.

It is not difficult to show that $C(m, a) \leq 1 + |a| = C(1, a)$, $m \in \mathbb{N}$. Therefore, the following question naturally arise.

Q1. *Is the sequence $C(m, a)$ monotone decreasing in m ?*

A partial answer is given in the next

Theorem 3.4. *The sequence $C(2^n, a), n \in \mathbb{N}$ is monotone decreasing in n .*

Proof. Indeed, since

$$\begin{aligned} \frac{|h^{2^{n+1}}(z) - h^{2^{n+1}}(w)|}{1 - \max\{|h(z)|^{2^{n+1}}, |h(w)|^{2^{n+1}}\}} &= \frac{|h^{2^n}(z) + h^{2^n}(w)|}{1 + \max\{|h(z)|^{2^n}, |h(w)|^{2^n}\}} \frac{|h^{2^n}(z) - h^{2^n}(w)|}{1 - \max\{|h(z)|^{2^n}, |h(w)|^{2^n}\}} \\ &\leq \frac{2 \max\{|h|^{2^n}(z), |h|^{2^n}(w)\}}{1 + \max\{|h(z)|^{2^n}, |h(w)|^{2^n}\}} \frac{|h^{2^n}(z) - h^{2^n}(w)|}{1 - \max\{|h(z)|^{2^n}, |h(w)|^{2^n}\}} \leq \frac{|h^{2^n}(z) - h^{2^n}(w)|}{1 - \max\{|h(z)|^{2^n}, |h(w)|^{2^n}\}}, \end{aligned}$$

we conclude that

$$j(h^{2^{n+1}}(z), h^{2^{n+1}}(w)) \leq j(h^{2^n}(z), h^{2^n}(w)),$$

i.e.,

$$C(2^{n+1}, a) = \sup_{z, w \in \mathbb{D}} \frac{j(h^{2^{n+1}}(z), h^{2^{n+1}}(w))}{j(z, w)} \leq \sup_{z, w \in \mathbb{D}} \frac{j(h^{2^n}(z), h^{2^n}(w))}{j(z, w)} = C(2^n, a).$$

\square

Q2. *Is it true that $C(m, (m+1)^{-1}) = 1$ for $m \geq 2$?*

Acknowledgement. The research of the second author was supported by the Academy of Finland grant with the Project number 2600066611.

REFERENCES

- [AVV] G.D. ANDERSON, M.K. VAMANAMURTHY, AND M. VUORINEN, *Conformal Invariants, Inequalities and Quasiconformal Maps*, John Wiley & Sons, New York, 1997.
- [AVZ] G.D. ANDERSON, M. VUORINEN, AND X. ZHANG, *Topics in special functions III*. - Manuscript 41 pp, arxiv.org/abs/1209.1696 [math.CA]
- [CCQ] M. CHEN, X. CHEN AND T. QIAN, Quasihyperbolic distance in punctured planes. *Complex Analysis and Operator Theory* (2012). In press/on-line DOI 10.1007/s11785-012-0275-1.
- [GO] F.W. GEHRING AND B.G. OSGOOD, Uniform domains and the quasihyperbolic metric, *J. Analyse Math.* 36, 1979, 50-74.
- [GP] F.W. GEHRING AND B.P. PALKA, Quasiconformally homogeneous domains, *J. Analyse Math.* 30, 1976, 172-199.
- [HIMPS] P. HÄSTÖ, Z. IBRAGIMOV, D. MINDA, S. PONNUSAMY AND S. K. SAHOO, Isometries of some hyperbolic-type path metrics, and the hyperbolic medial axis, In the tradition of Ahlfors-Bers, IV, *Contemporary Math.* 432, 2007, 63-74.
- [HPWW] M. HUANG, S. PONNUSAMY, H. WANG, AND X. WANG, A cosine inequality in the hyperbolic geometry. *Appl. Math. Lett.* 23 (2010), no. 8, 887-891.
- [K] R. KLÉN, Local convexity properties of quasihyperbolic balls in punctured space. *J. Math. Anal. Appl.* 342 (2008), no. 1, 192-201.
- [KVZ] R. KLÉN, M. VUORINEN, AND X. ZHANG, Quasihyperbolic metric and Möbius transformations. *Proc. Amer. Math. Soc.* (to appear) Manuscript 9pp, [arXiv: 1108.2967](https://arxiv.org/abs/1108.2967) math.CV.
- [RT1] A. RASILA AND J. TALPONEN, Convexity properties of quasihyperbolic balls on Banach spaces, *Ann. Acad. Sci. Fenn. Math.* 37, 2012, 215-228.
- [RT2] A. RASILA AND J. TALPONEN, On Quasihyperbolic Geodesics in Banach Spaces, [arXiv:1301.0900](https://arxiv.org/abs/1301.0900) [math.CV]
- [SVW] S. SIMIĆ, M. VUORINEN, AND G. WANG, Sharp Lipschitz constants for the distance ratio metric.- *Math. Scand.* (to appear), [arXiv:1202.6565](https://arxiv.org/abs/1202.6565) math.CA.
- [V] J. VÄISÄLÄ, The free quasiworld. Freely quasiconformal and related maps in Banach spaces. *Quasiconformal geometry and dynamics* (Lublin, 1996), Banach Center Publ., Vol. 48, Polish Academy of Science, Warsaw, ed. by Bogdan Bojarski, Julian Lawrynowicz, Olli Martio, Matti Vuorinen and Józef Zając, 1999, 55-118.
- [Vu1] M. VUORINEN, Conformal invariants and quasiregular mappings, *J. Analyse Math.*, **45** (1985), 69-115.
- [Vu2] M. VUORINEN, Conformal geometry and quasiregular mappings. *Lecture Notes in Math.* 1319, Springer-Verlag, Berlin-Heidelberg, 1988.

MATHEMATICAL INSTITUTE SANU, KNEZA MIHAILA 36, 11000 BELGRADE, SERBIA

E-mail address: `ssimic@turing.mi.sanu.ac.rs`

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF TURKU, 20014 TURKU, FINLAND

E-mail address: `vuorinen@utu.fi`